

Estimation of Regional-Global Methane Emissions and Refinement of Its Estimate by GOSAT-2 and Surface Observations

Principal Investigator: Naoko Saitoh

Institution: Chiba University

1-33, Yayoi-cho, Inage-ku, Chiba-shi, Chiba, JAPAN

Tel: +81-43-290-3843 / Fax: +81-43-290-3843

E-mail: naoko@faculty.chiba-u.jp

Cooperated by: Nara Women's University, Japan Agency for Marine-earth Science and Technology, Tokyo Gakugei University, and National Institute for Environmental Studies

[Abstract]

Key Words: Biogeochemical cycle, Global warming, Remote sensing, Methane, GOSAT/GOSAT-2, Surface observation, Chemical transport model

Methane (CH₄) is the second major anthropogenically produced greenhouse gas, after carbon dioxide. The objective of this study is to improve the accuracy of regional-global CH₄ emissions estimation through “top-down approach” using a chemical transport model, MIROC4-ACTM, and atmospheric observations. We retrieved CH₄ profiles from GOSAT, and estimated retrieval accuracy of 1% throughout the troposphere through comparisons with aircraft measurements and South Asian ground-based observations from our study. After validating the vertical and horizontal transport processes of MIROC4-ACTM through comparisons with aircraft and GOSAT measurements, we performed inversion analysis to estimate total CH₄ and nitrous oxide emissions. The inversion results are used in the IPCC 6th assessment report and could support the first global stocktake (GST). We also developed a LETKF data assimilation system for CH₄ flux estimation on model grid scale, and demonstrated that it was capable of reproducing known CH₄ flux patterns by assimilating pseudo-observations. We applied a two-box model simulation of carbon isotope of CH₄, $\delta^{13}\text{C-CH}_4$, for both hemispheres, and evaluated the consistency of global emission trends based on *a priori* and simulated *a posteriori* emissions with observed $\delta^{13}\text{C-CH}_4$ timeseries; the results suggested overestimation of CH₄ fugitive emissions in the EDGAR v4.3.2 inventory.

To better understand various types of regional CH₄ sources, particularly in South Asia, we have conducted long-term surface observations of CH₄, carbon monoxide (CO), and $\delta^{13}\text{C-CH}_4$ at Nainital, India and Comilla, Bangladesh. Simultaneous measurements of CH₄ atmospheric concentrations and fluxes from rice paddies in Tamil Nadu, India were also conducted. We evaluated the relationship between soil microbiology and greenhouse gases emissions quantitatively, and proposed a methodology for estimating global CH₄ emissions from rice paddies using information on soil microbial properties. The simultaneous measurements of concentrations and fluxes suggested that fluctuations in CH₄ emissions from rice paddies were an important factor in the seasonal variation of atmospheric CH₄ concentrations at the rice cultivation area. Diurnal variations in atmospheric CH₄ concentrations were less affected by CH₄ emissions from rice paddies than atmospheric transport in a one-dimensional advection model. Our trajectory analysis showed that most air masses with high CH₄ concentrations in northern India (Karnal and Sonapat) originated from the northwest, suggesting CH₄ sources in northwestern India. Analysis of the observed CH₄, CO, and $\delta^{13}\text{C-CH}_4$ timeseries at Comilla showed that 73% of CH₄ sources were of microbial origin and approximately 15% originated from biomass burning in autumn. The proportion of biomass burning sources increased to 32% in

winter.

[References]

- 1) P. K. PATRA, E. J. DLUGOKENCKY, J. W. ELKINS, G. S. DUTTON, Y. TOHJIMA, M. SASAKAWA, A. ITO, R. F. WEISS, M. MANIZZA, P. B. KRUMMEL, R. G. PRINN, S. O'DOHERTY, D. BIANCHI, C. NEVISON, E. SOLAZZO, H. LEE, S. JOO, E. A. KORT, S. MAITY, M. TAKIGAWA: *J. Meteorol. Soc. Japan, Ser. II*, 100, 2, in press, <https://doi.org/10.2151/jmsj.2022-018> (2022) (IF: 2.2)
Forward and inverse modelling of atmospheric nitrous oxide using MIROC4-atmospheric chemistry-transport model.
- 2) D. BELIKOV, N. SAITOH, P. K. PATRA, N. CHANDRA: *Remote Sensing*, 13(9), 1677 (2021) (IF: 4.848)
GOSAT CH₄ vertical profiles over the Indian subcontinent: effect of a priori and averaging kernels for climate applications.
- 3) J. S. H. BISHT, T. MACHIDA, N. CHANDRA, K. TSUBOI, P. K. PATRA, T. UMEZAWA, Y. NIWA, Y. SAWA, S. MORIMOTO, T. NAKAZAWA, N. SAITOH, M. TAKIGAWA: *J. Geophys. Res.*, 126, 4, e2020JD033541, <https://doi.org/10.1029/2020JD033541> (2021) (IF: 4.3)
Seasonal variations of SF₆, CO₂, CH₄ and N₂O in the UT/LS region due to emissions, transport and chemistry.
- 4) N. CHANDRA, P. K. PATRA, J. S. H. BISHT, A. ITO, T. UMEZAWA, N. SAIGUSA, S. MORIMOTO, S. AOKI, G. JANSSENS-MAENHOUT, R. FUJITA, M. TAKIGAWA, S. WATANABE, N. SAITOH, J. G. CANADELL: *J. Meteorol. Soc. Japan. Ser. II*, 99, 2309-2337, <https://doi.org/10.2151/jmsj.2021-015> (2021) (IF: 2.2)
Emissions from the Oil and Gas Sectors, Coal Mining and Ruminant Farming Drive CH₄ Growth over the Past Three Decades.
- 5) S. NOMURA, M. NAJA, M. K. AHMED, H. MUKAI, Y. TERAQ, T. MACHIDA, M. SASAKAWA, P. K. PATRA: *Atmos. Chem. Phys.*, 21, 16427-16452 (2021) (IF 6.133)
Measurement report: Regional characteristics of seasonal and long-term variations in greenhouse gases at Nainital, India, and Comilla, Bangladesh.
- 6) A. Z. OO, S. SUDO, T. FUMOTO, K. INUBUSHI, K. ONO, A. YAMAMOTO, S. D. BELLINGRATH-KIMURA, K. T. WIN, C. UMAMAGESWARI, K. S. BAMA, M. RAJU, K. VANITHA, P. ELAYAKUMAR, V. RAVI, V. AMBETHGAR: *Agriculture*, 10, 355, p16, doi:10.3390/agriculture10080355 (2020) (IF: 2.925)
Field validation of the DNDC-Rice model for CH₄ and nitrous oxide emissions from double-cropping paddy rice under different irrigation practices in Tamil Nadu, India.
- 7) A. Z. OO, S. SUDO, K. INUBUSHI, U. CHELLAPPAN, A. YAMAMOTO, K. ONO, M. MANO, S. HAYASHIDA, V. KOOTHAN, T. OSAWA, Y. TERAQ, J. PALANISAMY, E. PALANISAMY, R. VENKATACHALAM: *Agronomy*, 8, 202, <https://doi.org/10.3390/agronomy8100202> (2018) (IF: 3.417)
Mitigation Potential and Yield-Scaled Global Warming Potential of Early-Season Drainage from a Rice Paddy in Tamil Nadu, India.